

Reply to “Comment on ‘Tunnel Window’s Imprint on Dipolar Field Distributions’ ”

Stamp and Tupitsyn (ST) have recently posted a Comment [1] on a paper of ours [2]. They claim [1] that, in Ref. [2], “Short-time relaxation was caused by a simple noise field, acting uniformly over an ‘energy window’ of width $2\delta h_{hf}$ (to simulate the fluctuating nuclear spin bias)”. We see no basis for this statement.

Whereas previous work of Prokof’ev and Stamp [3, 4] deals with nuclear spin bath mechanisms that give rise to a tunnel energy window (TEW) for single-molecule magnets, such as Fe_8 , the work we report in Ref. [2] is independent of such mechanisms. The only definition of the TEW, δh_{hf} , we make use of in Ref. [2] is the one we give therein in the abstract: “we allow spin flips only if the corresponding energy change is less than some $2\delta h_{hf}$ ”. This is further explained below Eq. (2) in Ref. [2], where the system under consideration is defined as an Ising model of spins (that clearly stand for the electronic cluster spins) interacting through dipolar fields with the flipping rule just mentioned. Thus, the definition of δh_{hf} stands on its own without need for reference to the nuclear bath.

In addition, we mention hyperfine interactions only in passing, in an introductory remark, but the phrase “typical hyperfine coupling”, which PT ascribe to us, does not appear in Ref. [2]. We do refer once to δh_{hf} as “a typical hyperfine energy”, but make no use whatsoever of this non-definition.

We can similarly see no basis for the statement towards the end of the Comment by ST [1]: “... but here using the kinetic equation of Ref. [2], rather than just a phenomenological noise.”, since the Monte Carlo simulations we perform in Ref. [2] happen to be [5] a stochastic implementation of “the kinetic equation”, i.e., Eq. (4) of Ref. [3].

Finally, at the risk of being repetitious, we cannot understand the closing remark in the preceeding Comment, “...so we predict a steep fall of ξ_0 , which would be very hard to understand in the approach of ref. [4]”. Indeed, there is nothing to understand about this in the approach of Ref. [2], because the physics of nuclear baths which is behind TEW’s is not at all within the scope of Ref. [2].

What we do show in Ref. [2] is that a hole of half-width W develops in the dipole field distribution of an Ising model of spins interacting through dipolar fields with the flipping rule just mentioned, and that $W \simeq 0.75\delta h_{hf}$, and we know of no previously published work establishing a relation between these two quantities.

Juan J. Alonso¹ and Julio F. Fernández²

¹Física Aplicada I, Universidad de Málaga, 29071-Málaga, Spain. ²ICMA, CSIC and Universidad de Zaragoza, 50009-Zaragoza, Spain

-
- [1] P.C.E. Stamp and I. S. Tupitsyn, arXiv:cond-mat/0211413.
 - [2] J.J. Alonso and J.F. Fernández, Phys. Rev. Lett. **87**, 097205 (2001).
 - [3] N.V. Prokof’ev and P.C.E. Stamp, Phys. Rev. Lett., **80**, 5794 (1998).
 - [4] N.V. Prokof’ev and P.C.E. Stamp, J. Low Temp. Phys. **104**, 143 (1996).
 - [5] Except that we use a square hat like $\tau^{-1}(\xi)$ instead of an exponential function, and that Eq. (4) of Ref. [3] only holds for in the $\delta h_{hf}/k_B T \rightarrow 0$ limit, in the notation of Ref. [2].